

Journal Articles

- [1] C. Feuchter and H. Reinhardt, “Variational solution of the Yang–Mills Schrödinger equation in Coulomb gauge,” *Physical Review D - Particles, Fields, Gravitation and Cosmology*, 70, 105021, 2004. DOI: [10.1103/PhysRevD.70.105021](https://doi.org/10.1103/PhysRevD.70.105021).
- [2] C. Feuchter and H. Reinhardt, “Quark and gluon confinement in Yang–Mills theory in Coulomb gauge with the variational approach,” *Nuclear Physics B - Proceedings Supplements*, 141, 205–210, 2005. DOI: [10.1016/j.nuclphysbps.2004.12.031](https://doi.org/10.1016/j.nuclphysbps.2004.12.031).
- [3] D. Epple, C. Feuchter, and H. Reinhardt, “Quark and gluon confinement within the variational approach to Yang–Mills theory in Coulomb gauge,” *Nuclear Physics B - Proceedings Supplements*, 153, 234–241, 2006. DOI: [10.1016/j.nuclphysbps.2006.01.041](https://doi.org/10.1016/j.nuclphysbps.2006.01.041).
- [4] C. Feuchter and H. Reinhardt, “Yang–Mills vacuum in Coulomb gauge in $D = 2 + 1$ dimensions,” *Physical Review D - Particles, Fields, Gravitation and Cosmology*, 77, 085023, 2008. DOI: [10.1103/PhysRevD.77.085023](https://doi.org/10.1103/PhysRevD.77.085023).
- [5] G. Bälz, C. Feuchter, R. Handel, and B. Renz, “Filter media pore space analysis based on geometrical characteristics,” *Chemical Engineering and Technology*, 39, 499–504, 2016. DOI: [10.1002/ceat.201500348](https://doi.org/10.1002/ceat.201500348).
- [6] C. Feuchter and W. Schleifenbaum, “High-order lattice Boltzmann models for wall-bounded flows at finite Knudsen numbers,” *Physical Review E*, 94, 013304, 2016. DOI: [10.1103/PhysRevE.94.013304](https://doi.org/10.1103/PhysRevE.94.013304).
- [7] C. Feuchter, O. Wagner, A. Stief, and T. Beisswenger, “Turbulent flow simulations around a surface-mounted finite cylinder using an entropic multi-relaxation lattice Boltzmann method,” *Fluid Dynamics Research*, 51, 055509, 2019. DOI: [10.1088/1873-7005/ab3baf](https://doi.org/10.1088/1873-7005/ab3baf).
- [8] C. Feuchter, “Direct aeroacoustic simulation with a cumulant lattice-Boltzmann model,” *Computers and Fluids*, 224, 104970, 2021. DOI: [10.1016/j.compfluid.2021.104970](https://doi.org/10.1016/j.compfluid.2021.104970).

ArXiv Papers

- [1] D. Epple, C. Feuchter, and H. Reinhardt, “The Yang–Mills vacuum in Coulomb gauge,” Dec. 20, 2004. arXiv: [hep-th/0412231](https://arxiv.org/abs/hep-th/0412231).
- [2] C. Feuchter and H. Reinhardt, “Quark and gluon confinement in Coulomb gauge,” Feb. 13, 2004. arXiv: [hep-th/0402106](https://arxiv.org/abs/hep-th/0402106).
- [3] C. Feuchter and H. Reinhardt, “Variational solution of the Yang–Mills Schrödinger equation in Coulomb gauge,” Aug. 31, 2004. arXiv: [hep-th/0408236](https://arxiv.org/abs/hep-th/0408236).
- [4] H. Reinhardt and C. Feuchter, “On the Yang–Mills wave functional in Coulomb gauge,” Aug. 31, 2004. arXiv: [hep-th/0408237](https://arxiv.org/abs/hep-th/0408237).
- [5] C. Feuchter and H. Reinhardt, “The Yang–Mills vacuum in Coulomb gauge in $D = 2 + 1$ dimensions,” Nov. 15, 2007. arXiv: [0711.2452 \[hep-th\]](https://arxiv.org/abs/0711.2452).
- [6] H. Reinhardt, W. Schleifenbaum, D. Epple, and C. Feuchter, “Hamiltonian approach to Coulomb gauge Yang–Mills theory,” Oct. 1, 2007. arXiv: [0710.0316 \[hep-th\]](https://arxiv.org/abs/0710.0316).
- [7] H. Reinhardt, G. Burgio, D. R. Campagnari, D. Epple, C. Feuchter, M. Leder, M. Pak, J. M. Pawłowski, M. Quandt, W. Schleifenbaum, and A. Weber, “Hamiltonian approach to Yang–Mills theory in Coulomb gauge,” Nov. 3, 2009. arXiv: [0911.0613 \[hep-th\]](https://arxiv.org/abs/0911.0613).

Conference Papers

- [1] D. Epple, C. Feuchter, and H. Reinhardt, “The Yang-Mills vacuum in Coulomb gauge,” *AIP Conference Proceedings*, (Villasimius, Italy), 756, 2005, 269–271. DOI: [10.1063/1.1920963](https://doi.org/10.1063/1.1920963).
- [2] H. Reinhardt, W. Schleifenbaum, D. Epple, and C. Feuchter, “Hamiltonian approach to Coulomb gauge Yang-Mills theory,” *Proceedings of Science*, (Regensburg, Germany), 42, 2008. DOI: [10.22323/1.042.0326](https://doi.org/10.22323/1.042.0326).
- [3] H. Reinhardt, G. Burgio, D. R. Campagnari, D. Epple, C. Feuchter, M. Leder, M. Pak, J. M. Pawlowski, M. Quandt, W. Schleifenbaum, and A. Weber, “Hamiltonian approach to Yang-Mills theory in Coulomb gauge,” *Proceedings of Science*, (Trento, Italy), 87, 2010. DOI: [10.22323/1.087.0038](https://doi.org/10.22323/1.087.0038).
- [4] C. Feuchter and A. Schneider, “Direct aeroacoustics simulations using modern lattice-Boltzmann methods,” *Fortschritte der Akustik DAGA 2024*, (Hanover, Germany), 2024. https://pub.dega-akustik.de/DAGA_2024/konferenz?article=332.